

DTPA • Chelated Micronutrient for Plant Nutrition

ChelStar Iron 11 • 7-0-0 GUARANTEED ANALYSIS

Total Nitrogen (N)......7.0% 7.0% Water-Soluble Nitrogen Iron (Fe).....11.0%

11.0% Chelated Iron

Derived from iron diethylenetriaminepentaacetate.

Precautionary Statements: This product is not classified as a hazardous substance based on currently available data. However, observe good industrial hygiene practices. Wash hands after handling.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Keep out of reach of children. Store away from incompatible materials. Dispose of contents/container in accordance with local authority regulations.

GENERAL INFORMATION:

ChelStar Iron 11 is an DTPA chelated, water-soluble, powdered form of iron, useful for prevention and correction of iron deficiency in crops. Soil application is the preferred application method to prevent iron deficiency. Foliar application, however, will provide correction when a soil application is impractical.

SOIL APPLICATION DIRECTIONS:

Apply *ChelStar Iron 11* to acid and mildly alkaline soils. It should be dissolved in water or fluid fertilizer, or dry-blended with water-soluble fertilizers for optimum delivery and coverage. For trees or individual plants, *ChelStar Iron 11* may be blended with an inert (such as soil or sand) or sprinkled directly on the soil uniformly under the plant's drip line, then watered in. Soil applications may also be made by dissolving in water and then mixing or metering into drip, sprinkler, or furrow irrigation systems.

FOLIAR APPLICATION DIRECTIONS:

ChelStar Iron 11 may be applied in water or in combination with most pesticides. Buffer the spray solution to a pH of 6 to 6.5 for best performance. Thorough coverage and the use of wetting agents often enhance nutrient uptake from foliar sprays. Application rates and dilution factors depend on crop sensitivity, the amount of foliage to be sprayed, and the application method. Avoid applying ChelStar Iron 11 when plants are suffering from moisture stress. If there is any doubt, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity before general application.

AERIAL APPLICATION DIRECTIONS: Apply 0.33 lb. of *ChelStar Iron 11* per acre in at least 10 gal. of water. For application timing and frequency, refer to appropriate foliar application recommendation below.

CROP RECOMMENDATIONS – SOIL APPLICATION

VEGETABLE CROPS (including beans, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, sweet corn, tomatoes) and FIELD CROPS (including alfalfa, corn, dry beans, peas, forage grasses, rice, safflower, sorghum, sugar beets, sugarcane): Apply ChelStar Iron 11 at 4 to 12 lbs. per acre.

TREE & VINE CROPS (including almond, apple, apricot, avocado, berry, cherry, citrus, grape, mango, olive, pear, peach, pistachio, plum, walnut): Apply ChelStar Iron 11 at 4 to 15 lbs. per acre or 1 to 4 oz. per inch of tree diameter.

ORNAMENTAL SHRUBS & TREES (including azaleas, gardenias, junipers, pines, roses): Apply *ChelStar Iron 11* at 2 Tbsp. for small shrubs and up to 2 Tbsp. per inch of tree diameter for large trees and shrubs.

TURF: Apply *ChelStar Iron 6* at 0.75 lb. per 1,000 sq. ft. or 33 lbs. per acre.

Note: the rates provided above are based on broadcast application. If broadcast rates are used for spot or banded applications, phytotoxicity may result.

CROP RECOMMENDATIONS – FOLIAR APPLICATION

TREES (CITRUS): Apply *ChelStar Iron 11* at 1 to 2 lbs. per 100 gal. of cover spray any time except during bloom.

TREES & VINE CROPS (excluding citrus): Apply ChelStar Iron 11 at 1 to 2 lbs. per acre in sufficient spray volume for thorough coverage (25 gal. minimum) at dormant/delayed dormant before bloom, then after full bloom at 2 to 3 week intervals as needed. NOTE: For smoothskinned fruit (peach, pear, apple, etc.) apply 1lb. per acre in 50 to 100 gal. of water.

VEGETABLE & FIELD CROPS (ground equipment only): Apply *ChelStar Iron 11* at 0.33 to 1.33 lbs. per acre in sufficient spray volume for thorough coverage (25 gal. minimum).

TURF: Apply *ChelStar Iron 11* at 1 to 4 oz. per 1,000 sq. ft. With growth regulators, apply 1 to 3 oz. per 1,000 sq. ft. Always apply in sufficient spray volume for thorough coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.html

WARNING: This product contains a chemical known to the State of California to cause cancer and birth defects or other reproductive harm.

CONDITIONS OF SALE AND WARRANTY

Chemurgic Agricultural Chemicals, Inc. warrants that this product conforms to the chemical description on the label and is reasonably suited for the purposes stated when used as recommended under normal conditions.

Unintended consequences such as crop damage or poor performance may result from factors including, but not limited to, method of application, timing, crop and soil condition, weather, and other materials present, all of which are beyond the control of *Chemurgic Agricultural Chemicals, Inc.* and Seller.

Chemurgic Agricultural Chemicals, Inc. neither makes, nor authorizes any agent or representative to make any other warranties, expressed or implied, including fitness or merchantability.

Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material.

In no case shall *Chemurgic Agricultural Chemicals, Inc.* or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product.

The foregoing is a condition of sale by *Chemurgic Agricultural Chemicals, Inc.* and is accepted as such by the Buyer.



Chemurgic Agricultural Chemicals, Inc.

P.O. Box 2106 • Turlock, CA 95381

Net Weight: 25 Lbs (11.3 Kg)